

COMERCIAL NAME	COMPOSITION	TARGET SPECIES	ADMINISTRATION ROUTE AND POSOLOGY	PRES.
HORMONAL				
OXIVEEX Injectable solution	Oxytocin 10 IU Excipients q.s. 1 ml	Sows Cows Mares Ewes Does	Cows and Mares: Intramuscular or subcutaneous route: 3 to 5 ml. Slow intravenous route: 1.5 to 3 ml. Sows and ewes: Intramuscular or subcutaneous route: 2 to 3 ml. Slow intravenous route: 1 to 1.5 ml.	50 ml
TESTOSTERONA-E Injectable oily solution	Testosterone (propionate) 25 mg Vitamin E acetate 50 mg Excipients q.s. 1 ml	Bovine Equine Ovine Caprine	Intramuscular route Cattle and horses: 10 ml/animal. Pigs, sheep and goats: 2 ml/animal. Administer twice a week for 1 to 3 weeks.	50 ml
INTRAMAMMARY SUSPENSIONS				
MASTIVEEX CON CEFALOSPORINA Intramammary suspension	Ceftiofur hydrochloride 125 mg Dexamethasone (21-phosphate sodium).....2 mg Excipients q.s. 10 g	Bovine	One syringe per affected quarter every 24 hours for a minimum of 2 days.	10 g
MASTIVEEX DRY Intramammary suspension	Cloxacillin (benzathine) 500 mg Excipients q.s. 5 g	Bovine Ovine Caprine	Administer the entire content of the syringe in each quarter in a single dose after the last milking.	5 g
MASTIVEEX LACTACIÓN Intramammary suspension	Cloxacillin (sodium) 200 mg Ampicillin (sodium) 75 mg Excipients q.s. 10 g	Bovine Ovine Caprine	One syringe in each affected quarter every 12 hours, for 3 consecutive milkings.	10 g 5 g
ORAL PHARMACEUTICAL FORMS				
POWDERS FOR ORAL ADMINISTRATION				
AMOXICILINA 50% REVEEX Oral powder	Amoxicillin (trihydrate) 500 mg Excipients q.s. 1 g	Porcine Poultry	Oral administration through drinking water Broilers: 15 mg/kg b.w./day, for 5 days Ducks for fattening: 20 mg/kg b.w./day for 3 days Turkeys for fattening: 15 to 20 mg/kg b.w./day for 5 days Pigs: 20 mg/kg b.w./day for 4 days	200 g 1 kg 25 x 1 kg
COLISTINA REVEEX 2050000 UI/g Oral powder	Colistin sulphate 2050000 IU Excipients q.s. 1 g	Bovine Porcine Poultry Turkeys	Oral administration through drinking water Calves and pigs: 0.5 g/10 kg b.w./day for 3 days Poultry and turkeys: 0.06 g/kg b.w./ day for 3 days	200 g 1 kg 25 x 1 kg
DOXICICLINA 50 Oral powder	Doxycycline (hyclate) 500 mg Excipients q.s. 1 g	Bovine Porcine Poultry	Oral administration through drinking water Poultry: 30 mg/kg b.w./day for 3 to 5 days Fattening pigs: 0.2 g/10 kg b.w./day for 5 days Calves: 0.2 g/10 kg b.w./day for 5 days	200 g 1 kg 25 x 1 kg
LINCOMICINA 40 Oral powder	Lincomycin (hydrochloride) 400 mg Excipients q.s. 1 g	Porcine Poultry	Oral administration through drinking water Pigs: 12.5 - 25 mg/kg b.w./day for 5 to 10 days. Poultry: Administer 7.5-15 mg/kg b.w./day, for 7 days.	100 g 1 kg
TYLOSINA 500 REVEEX Oral powder	Tylosin (tartrate) 500 mg Excipients q.s. 1 g	Bovine Porcine Poultry	Oral administration through drinking water Calves: 20-40 mg/kg b.w. every 12 hours for 7 to 14 days Pigs: 40 mg/kg b.w. for 10 days Poultry: 150-200 mg/kg b.w. for 3 to 5 days	100 g 250 g 500 g 1 kg
SOLUTIONS AND SUSPENSIONS FOR ORAL ADMINISTRATION				
ALBENDEX 10% Oral suspension	Albendazole 100 mg Excipients q.s. 1 ml	Bovine Ovine Caprine	Oral route Gastrointestinal nematodes, strongylida lungworms and cestodes: 7.5 ml/100 kg b.w. in a single dose Adult flukes: 10 ml/100 kg b.w. in a single dose	200 ml 1 L 1 gallon
AMPROLIUM 40% REVEEX Oral solution	Amprolium (hydrochloride) 400 mg Excipients q.s. 1 ml	Poultry	Oral administration through drinking water 0.5 ml/10 kg b.w./day for 5-7 days	500 ml 1 L 5 L
BRONCOVEEX Oral solution	Bromhexine (hydrochloride) 10 mg Excipients q.s. 1 ml	Bovine Porcine Caprine Equine Poultry Ovine	Oral administration through drinking water Poultry: 100 ml in 1000 L of water for 5 days Pigs: 0.2 to 0.5 ml/animal for 5 days Piglets: 0.1 to 0.25 ml/animal for 5 days Calves: 0.3 to 0.7 ml/animal for 5 days Cattle and horses: 1 to 1.5 ml/animal for 5 days	1 L 1 gallon
CARNIVIT AMINO Oral solution	Vitamin B1 hydrochloride 2.85 mg; Vitamin B2-5-sodium phosphate 3.2 mg; Vitamin B6 hydrochloride 1.8 mg; Biotin 0.05 mg; Nicotinamide 10 mg; Dexpanthenol 5 mg; Vitamin A 325 IU; Vitamin D3 100 IU; Vitamin 12.5 mg; Vitamin E acetate 4.5 mg; L-Carnitine hydrochloride 12.5 mg; Betaine 25 mg; L-Arginine 3.75 mg; L-Glycine 7.5 mg; L-Lysine hydrochloride 12.9 mg; DL-Methionine 5.5 mg; L-Tryptophan 1.8 mg; L-Threonine 3.2 mg; Excipients q.s. 1 ml	Cattle Horses Pigs Poultry Sheep Goats Rabbits	Oral, in drinking water. All species: 2 ml/L of water for 5 consecutive days	100 ml 1 L 18 L
COCXIVET Oral solution	Toltrazuril 25 mg Excipients q.s. 1 ml	Poultry	Oral administration through drinking water 1 ml/l of drinking water for 2 days	100 ml 1 L 1 gallon
DICLAZURIL 2,5% Oral solution	Diclazuril 25 mg Excipients q.s. 1 ml	Bovine Ovine Poultry Rabbits	Oral administration through drinking water Cattle, sheep, goats: 1 ml/25 kg b.w. in a single dose Rabbits: 0.002 ml/kg b.w. in a single dose Poultry: 0.3 mg/kg b.w./day for 3 days	200 ml 1.5 L 1 gallon

COMERCIAL NAME	COMPOSITION	TARGET SPECIES	ADMINISTRATION ROUTE AND POSOLOGY	PRES.
DOXYVEEX 200 MG/ML Oral solution	Doxicycline (hyclate) 200 mg Excipients q.s. 1 ml	Porcine Poultry Turkeys	Oral administration through drinking water Porcine: 1 ml/20 kg b.w. for 5 days Poultry and turkeys: 0.05-0.1 ml/kg b.w. for 3 to 5 days	1 L 1 gallon 5 L
ENROFLOXACINA 20% REVEEX Oral solution	Enrofloxacin 200 mg Excipients q.s. 1 ml	Poultry	Oral administration through drinking water 0.05 ml/kg b.w./day per day for 3 to 5 days.	250 ml 1 L 1 gallon
FENIVEEX Oral solution	Florfenicol 100 mg Excipients q.s. 1 ml	Porcine Poultry	Oral administration through drinking water 1 ml/l drinking water/day for 3 to 5 days	1 L 1 gallon
HEPATOVEEX SOLUCIÓN ORAL Oral solution	Lysine hydrochloride 20 mg Betaine 60 mg Choline chloride 75 mg DL-Methionine 11.5 mg Sorbitol 350 mg Artichoke extract.....10 mg Taurine 12.5 mg Carnitine Hydrochloride 25 mg Vitamin B12 10 µg Excipients q.s 1 ml	Poultry Pigs Cattle Sheep Goats Camelids Horses Rabbits	Oral, in drinking water. Poultry and pigs: 1 ml/L of water for 5 consecutive days. Cattle and horses: 1 ml/5-10 kg b.w. for 5 consecutive days Sheep, goats, camelids, rabbits: 1 ml/2.5-5 kg b.w. for 5 consecutive days.	1 L 5 L 25 L
HIDRO REX VITAL AMINOACIDOS Oral solution	Vitamin B1 5 mg; Vitamin B2 10 mg; Vitamin B6 3 mg; Vitamin B12 30 µg; Nicotinic acid 20 mg; Calcium pantothenate 10 mg; Vitamin C 50 mg; Vitamin K3 5 mg; Folic acid 1 mg; Vitamin A 20,000 IU; Vitamin D3 5,000 IU; Vitamin E 9 mg; Aspartic acid 14.5 mg; Threonine 6.4 mg; Serine 6.6 mg; Glutamic acid 26.4 mg; Proline 11.5 mg; Glycine 15.3 mg; Alanine 17.4 mg; Cystine 1.26 mg; Lysine 20.7 mg; Histidine 5.6 mg; Arginine 14.1 mg; Methionine 11.1 mg; Tryptophan 3.67 mg; Isoleucine 9.8 mg; Leucine 20.1 mg; Phenylalanine 7.7 mg; Tyrosine 6.8 mg; Excipients q.s. 1 ml	Bovine Equine Porcine Poultry Ovine Caprine Rabbits	Orally, in drinking water. Poultry: 0.5-1 ml/L water for 5 days Cattle: <i>Young animals:</i> 1 ml/L milk for 5 days <i>Adults:</i> 0.5-1 ml/L water for 5 days Pigs: <i>Piglets:</i> 2 ml/L water for 5 days <i>Fattening:</i> 0.5 ml/L water for 5 days	100 ml 1 L 18 L
PARACETAMOL LÍQUIDO 30% Oral solution	Paracetamol 300 mg Excipients q.s. 1 ml	Poultry	Oral administration through drinking water 1ml/10 kg b.w. for 5 days	1L 5 L 1 gallon
SULFATRIM Oral solution	Sulfamethoxypyridazine 200 mg Trimethoprim 40 mg Excipients q.s. 1 ml	Bovine Ovine Caprine Poultry	Oral administration through drinking water Bovine, ovine and caprine: 1 ml/L drinking water for 7 days Poultry: 0.4-0.8 ml/L drinking water for 3 to 5 days	1 L 5 L
VETIFOS Oral solution	Zinc (chloride monohydrate) 4 mg; Copper (chloride dihydrate) 3 mg; Manganese (chloride tetrahydrate) 2 mg; Calcium (chloride dihydrate) 23 mg; Magnesium (chloride hexahydrate) 4.5 mg; Phosphorus (monosodium phosphate) 100 mg; Citric acid 204mg; Excipients q.s. 1ml	Poultry Porcine Bovine Ovine Caprine	Oral administration through drinking water. Bovine, ovine, caprine and porcine: 10 to 15 ml/100 kg b.w./day for 5 to 10 days. Poultry: 5 to 10 ml/L of water for 5 to 10 days.	1 L 5 L

INTRAUTERINE TABLETS				
VEEX-CICLIN 1000 MG Intrauterine tablets	Chlortetracycline hydrochloride 1000 mg Excipients q.s. 8 g	Bovine	2 tablets per cow per day, in a single dose	40 × 8 g

BIOCIDES				
CIPERZURON Self-emulsifying concentrate	Cypermethrin 75 mg Diflubenzuron 45 mg Excipients q.s. 1 ml	--	Flying insects: 5 ml in 1 L of water (0.5%). Regular sanitization: 10 ml in 1 L of water (1%). Heavy infestations: 20 ml in 1 L of water (2%).	1 L 5 L
CLORHEX-VEEX ESPUMANTE Solution	Chlorhexidine (digluconate) 200 mg Excipients q.s. 1 ml	Porcine	Direct application on the skin of the animals, diluted to 5-10% in water, depending on the amount of dirt and wounds.	5 L 25 L
FENOLVEEX POULTRY Self-emulsifying concentrate	p-chloro-m-cresol 250 mg Salicylic acid 100 mg Excipients q.s. 1 ml	--	Routine disinfection: 2 L/100 L of water for 30 minutes. Coccidial oocysts: apply 3-4 L/100 L of water for 2 hours. Encapsulated viruses: use 1 L/100 L of water for 1 hour.	1 L 5 L
GLUTALVEEX QUATS Solution	Glutaraldehyde 150 mg Didecylidimethyl ammonium chloride 100 mg Contains citronella essential oil Excipients q.s 1 ml	--	3 ml/L of water to treat 10m². For disinfection of ASF virus administer 5 ml/L of water and apply 3 L of solution for every 10 m².	1 L 5 L
GLUTEX Solution	Glutaraldehyde 40 mg Gloxal 32 mg Didecylidimethyl ammonium chloride 100 mg Excipients q.s 1 ml	--	Spraying, sprinkling or washing at a rate of 1 L/200 L of water.	1 L 5 L 1 gallon
YODORVEEX Solution	Iodine 25 mg Excipients q.s. 1 ml	--	Floors, walls and equipment: 10 - 50 ml/l of water. Manual or automatic spraying: 25 ml product/m². Sanitary footbaths: 10 - 50 ml/L of water.	1 L 5 L

GENERAL
PORTFOLIO
.....

VETERINARY
MEDICINAL
PRODUCTS

REVEEX

Av. Antón Phillips, cruce con calle El Canal - Complejo Industrial Reveex - Zona Industrial La Hamaca
Tel. +58-243-551,54,32 /64,03/68,52/69,76 · Fax. +58-243-551,68,53 · Website: <https://reveex.com/laboratorios>
MARACAY - ESTADO ARAGUA - VENEZUELA

REVEEX Laboratories also has specific portfolios
for trypanocidal and aquaculture products

COMERCIAL NAME	COMPOSITION	TARGET SPECIES	ADMINISTRATION ROUTE AND POSOLOGY	PRES.
INJECTABLE DRUGS				
ENDOPARASITICIDES				
BOVINET Injectable solution	Ivermectin 31.5 mg Excipients q.s. 1 ml	Bovine Ovine Caprine	Subcutaneous route 1 ml/50 kg b.w. in a single dose	50 ml 100 ml 250 ml 500 ml
IVERMERCITINA 1% Injectable solution	Ivermectin 10 mg Excipients q.s. 1 ml	Bovine Ovine Caprine	Subcutaneous route 1 ml/50 kg b.w. in a single dose 1.5 ml/50 kg b.w. in a single dose	10 ml 50 ml 100 ml 500 ml
IVERMECTINA FS Solución inyectable	Ivermectina 10 mg Clorsulón 100 mg Excipientes csp 1 ml	Bovino	Vía subcutánea Bovinos: 1 ml/50 kg p.v. dosis única	50 ml 100 ml 500 ml

ANTIMICROBIALS				
DIHIDROESTREX 250 MG/ML Injectable solution	Dihydrostreptomicine (sulphate) 250 mg Excipients q.s. 1 ml	Bovine Porcine	Intramuscular route 1 ml/10 kg b.w./day for 3 to 4 days	100 ml 250 ml
DOXIVEEX 200 MG/ML Injectable solution	Doxycycline (hyclate) 200 mg Excipients q.s. 1 ml	Bovine Porcine	Intramuscular route 0.5 - 1 ml/10 kg b.w. for 4 to 5 days	100 ml 250 ml
FENIVEEX Injectable solution	Florfenicol 300 mg Excipients q.s. 1 ml	Bovine Porcine Ovine Caprine	Intramuscular route 1 ml/15 kg b.w. in 2 doses in 2 doses with an interval of 48 hours	20 ml 100 ml 250 ml
FLORFENIXIN Injectable solution	Florfenicol 300 mg Flunixin (meglumine) 16,5 mg Excipients q.s. 1 ml	Cattle	Subcutaneous route Administer 2 ml/15 kg b.w. in a single dose.	50 ml 100 ml 250 ml
GENTAVEX 12% Injectable solution	Gentamicin (sulphate) 120 mg Excipients q.s. 1 ml	Bovine Porcine Equine Canine	Intramuscular route Bovine, equine and porcine: 1 ml/40 mg b.w. every 12 hours for 3 days. Canine: 0.2/8 kg b.w. every 12 hours for 4 to 5 days	100 ml 250 ml
LINCOSIN-100 Injectable solution	Lincomycin (hydrochloride) 100 mg Excipients q.s. 1 ml	Porcine	Intramuscular route Pigs: 1.1 ml/10 kg b.w., for 3 to 7 days.	100 ml 250 ml
LINCOVEEX Injectable solution	Spectinomycin (sulfate) 100 mg Lincomycin (hydrochloride) 50 mg Excipients q.s. 1 ml	Bovine Porcine	Intramuscular route 1 ml/kg b.w./day for 3 to 5 days	100 ml 250 ml
MARBOFLOXACINA 10% Injectable solution	Marbofloxacin 100 mg Excipients q.s. 1 ml	Bovine Porcine Canine Feline	Intramuscular or subcutaneous route Cattle: 2 ml/25 kg b.w. in a single dose Pigs: 1 ml/50 kg b.w. for 3-5 days.	20 ml 50 ml 100 ml 250 ml 500 ml
QUINOCOLI 10% Injectable solution	Enrofloxacin 100 mg Excipients q.s. 1 ml	Bovine Porcine	Intramuscular or subcutaneous route Cattle: 7.5 ml/100 kg b.w. per day Pigs: 0.75 ml/10 kg b.w. In severe cases repeat treatment after 48 hours.	20 ml 100 ml 250 ml
SEPTIBRON FORTE TF Injectable solution	Florfenicol 150 mg Tylosin (tartrate) 100 mg Excipients q.s. 1 ml	Bovine Porcine Ovine Caprine	Deep intramuscular injection. Cattle, sheep, goats and buffalo: Administer 1 ml/10 kg body weight. Repeat after 48 hours. Pigs: Administer 1 ml/30 kg body weight. Repeat after 48 hours.	20 ml 50 ml 100 ml 250 ml
TETRALONG 20 L.A. Injectable solution	Oxytetracycline (dihydrate).....200 mg Excipients q.s. 1 ml	Bovine Porcine Ovine Caprine	Deep intramuscular or subcutaneous route Bovine, porcine, ovine and caprine: 1 ml/10 kg b.w. In severe cases repeat treatment after 48-72 hours. Piglets: 1ml/piglet subcutaneously.	50 ml 100 ml 250 ml
TETRALONG 30 L.A. Injectable solution	Oxytetracycline (dihydrate) 300 mg Excipients q.s. 1 ml	Bovine Porcine	Deep intramuscular or subcutaneous route Bovine and porcine: 1 ml/10 kg b.w., in a single dose. In severe cases repeat treatment after 48-72 hours.	50 ml 100 ml
TULASOL Injectable solution	Tulathromycin 100 mg Excipients q.s. 1 ml	Bovine Porcine Ovine	Intramuscular or subcutaneous route Bovine, porcine and ovine: 1 ml/40 kg b.w. in a single dose.	50 ml 100 ml
TYLOSINA 200 Injectable solution	Tylosin (tartrate) 200 mg Excipients q.s. 1 ml	Bovine Porcine Ovine Caprine	Intramuscular route. Adult bovine: 0.5 to 1 ml/20 kg b.w./day. Calves, porcine, ovine and caprine: 1 ml/20 kg b.w./day. Piglets: 0.25 ml/5 kg b.w./day. Administer for no longer than 5 days.	100 ml 250 ml
ULTRADIAZINE Injectable solution	Sulfadiazine 200 mg Trimethoprim 40 mg Excipients q.s. 1 ml	Bovine Porcine Ovine Caprine	Intramuscular route. 1 ml/10 kg b.w./day for 2 to 3 days. In severe cases repeat after 12 hours.	100 ml 250 ml

COMERCIAL NAME	COMPOSITION	TARGET SPECIES	ADMINISTRATION ROUTE AND POSOLOGY	PRES.
BETA-LACTAMS				
15-AMOX L.A. Injectable oily suspension	Amoxicillin (trihydrate) 150 mg Excipients q.s. 1 ml	Bovine Porcine Ovine Caprine	Intramuscular or subcutaneous route. 1 ml/10 kg b.w., repeat after 48 hours.	20 ml 50 ml 100 ml
CEPTIFUR Injectable oily suspension	Ceftiofur (hydrochloride) 50 mg Excipients q.s. 1 ml	Bovine Porcine	Intramuscular or subcutaneous route Cattle: 1 ml/50 kg b.w. for 3 to 5 days. Pigs: 1 ml/16 kg b.w. for 3 days.	50 ml 100 ml 250 ml
ENROAMOX Injectable oily suspension	Amoxicillin (trihydrate) 100 mg Enrofloxacin 50 mg Excipients q.s. 1 ml	Bovine Porcine	Intramuscular or subcutaneous route 0.5 - 1 ml/10 kg b.w. for 3 to 5 days	50 ml 100 ml 250 ml
PENIVEEX-200 RETARD Injectable suspension	Benzy/penicillin procaine 100000 UI Benzathine benzylpenicillin 100000 UI Dihydrostreptomycin (sulfate) 200 mg Chlorphenamine maleate 1 mg Excipients q.s. 1 ml	Bovine Equine Porcine Ovine Caprine	Intramuscular or subcutaneous route. Adult cattle and horses, pigs, sheep and goats: 5-10 ml/100 kg b.w. Calves and piglets: 0.75 ml PENIVEEX/10 kg b.w. Administer 2 or 3 injections at 48-hour intervals.	100 ml 250 ml
PENIVEEX ESTREPTO 200 Injectable suspension	Benzy/penicillin procaine 200 mg Dihydrostreptomycin (sulfate) 200 mg Excipients q.s. 1 ml	Bovine Equine Porcine Ovine Caprine	Intramuscular or subcutaneous route. Adult cattle and horses, pigs, sheep and goats: 5-10 ml/100 kg b.w. Calves and piglets: 0.75 ml PENIVEEX/10 kg b.w. Administer 2 or 3 injections at 48-hour intervals.	100 ml 250 ml
PENIVEEX-300 Injectable suspension	Benzy/penicillin procaine 300 mg Excipients q.s. 1 ml	Bovine Porcine Equine Ovine	Intramuscular route Bovine and equine: 0.4 ml/10 kg b.w. for 3 days Ovine and porcine: 0.5 ml/10 kg b.w. for 3 days	100 ml 250 ml

VITAMINS AND MINERALS				
ADEX-3 FORTE Hydrodispersible injectable solution	Vitamin A 500000 IU Vitamin D3 75000 IU Vitamin E 50 mg Excipients q.s. 1 ml	Bovine Porcine Ovine Caprine	Intramuscular route Adult cattle: 6 - 8 ml/animal. Calves and adult pigs: 4 ml/animal. Fattening pigs, sheep and Caprines: 2 ml/animal. Piglets, lambs and kids: 1 ml/animal. Repeat treatment after 30 days if necessary.	20 ml 100 ml 250 ml 500 ml
AMINOVEEX INYECTABLE Injectable solution	Vitamin A 30000 UI Vitamin D3 10000 UI Vitamin E 10 mg Magnesium hypophosphite 5 mg Manganese sulphate 0.7 mg Copper sulphate 0.95 mg Magnesium chloride 1 mg Cobalt chloride 0.1 mg L-Arginine 20 mg Acetylmethionine 100 mg Excipients q.s. 1 ml	Bovine Porcine Equine Ovine Caprine	Intramuscular route. For horses and pigs, intramuscular injection only. For cattle, sheep and goats: intramuscular or subcutaneous injection. Cattle and horses: 2 ml per 100 kg body weight. Pigs, sheep and goats: 1 ml per 15 kg body weight. Repeat the dose every 15 days.	50 ml 100 ml 250 ml
CALCIOVEEX 2:2:1 Injectable solution	Calcium gluconate 155 mg Calcium hypophosphite 27.5 mg Magnesium chloride 83.5 mg Glucose 176 mg Excipients q.s. 1 ml	Bovine Equine Ovine Caprine	Slow intravenous, intramuscular or subcutaneous routes. Bovines and equines (500 kg b.w. approx.): 500 ml. Calves and foals: 50-100 ml. Ovines and caprines: 25-50 ml.	250 ml 500 ml
ESVEEX Injectable emulsion	Vitamin E 150 mg Selenium (sodium selenite) 0.5 mg Excipients q.s. 1 ml	Bovino Porcino Ovino	Intramuscular route Calves: 5 ml/animal, repeat after 7 days. Piglets: 0.5 ml/animal, repeat after 7 days. Lambs: 1 ml/animal, repeat after 7 days.	25 ml 50 ml 100 ml
FERTOL Injectable suspension	Iron (III) (complex with Dextran) 182 mg Toltrazuril 36,5 mg Suitable excipients q.s. 1 ml	Newborn piglets	Intramuscular route. 0.55 ml/kg b.w. (equivalent to 20 mg/kg b.w. of toltrazuril and 100 mg/kg b.w. of iron, as a single dose; should be administered to piglets between 48 and 72 hours after birth).	100 ml 250 ml
GLEPTOVEEX Injectable solution	Iron (III) as Gleptoferron complex 200 mg Excipients q.s. 1 ml	Porcine	Intramuscular route 1 ml/piglet within the first 3 days after birth	100 ml 250 ml
IRONVEEX 200 MG/ML Injectable solution	Iron (as Iron-Dextran) 200 mg Excipients q.s. 1 ml	Bovine Porcine Ovine	Deep intramuscular route Piglets: Curative: 1 ml/piglet and repeat after 7 days. Calves: Curative: 2.5 ml/calf and repeat after 7 days. Lambs: Curative: 1 ml/lamb and repeat after 7 days.	100 ml
NUTRIL VACAS LECHERAS Injectable solution	Calcium Hypophosphite 20 mg, Copper Chloride 1 mg, Cobalt Chloride 1 mg, Magnesium Chloride 1 mg, Potassium Iodide 3 mg, Sodium Selenite 1.5 mg, Vitamin B12 5 µg, Vitamin D2 1000 IU, Vitamin A 5000 IU, Excipients q.s. 1 ml	Bovine	Intramuscular route 1 ml per 30 kg b.w., every 90 days.	250 ml

COMERCIAL NAME	COMPOSITION	TARGET SPECIES	ADMINISTRATION ROUTE AND POSOLOGY	PRES.
NUTRIL Injectable solution	Vitamin A 5000 IU; Vitamin D2 20000 IU; Vitamin E 5 mg; Vitamin B12 50 µg; Vitamin B6 (hydrochloride) 8 mg; L-Tryptophan 0.25 mg; L-Leucine 2.1 mg; L-Lysine hydrochloride 10 mg; L-Valine 2 mg; L-Glycine 5 mg; L-Arginine hydrochloride 4 mg; DL-Methionine 4 mg; L-Histidine hydrochloride 2.1 mg; L-Threonine 0.25 mg; Copper sulfate 20 µg; Cobalt sulfate 0.13 mg; Zinc gluconate 0.10 mg; magnesium hypophosphite 2.1 mg; Calcium gluconate 15 mg; Sodium chloride 0.42 mg; Potassium iodide 0.15 mg; Sodium selenite 1 mg; Iron citrate (ammoniacal) 4 mg; Nandrolone decanotate 2 mg; Excipients q.s. 1 ml	Bovine Equine Ovine Caprine	Intramuscular or subcutaneous route Calves and Foals: 5 ml/animal. Adult cattle and horses: 10 ml/animal. Lambs and Kids: 2 - 3 ml/animal. Adult sheep and goat: 5 - 6 ml/animal.	100 ml 250 ml
OSMOLAR VITAMINADO Injectable solution	Glucose 50 mg; Calcium chloride 0.15 mg; Potassium chloride 0.25 mg; Magnesium hypophosphite 0.20 mg; Sodium acetate 2.5 mg; L-Histidine hydrochloride 20 µg; DL-Methionine 20 µg; L-Tryptophan 10 µg; L-Cysteine hydrochloride 10 µg; L-Threonine 35 µg; L-Isoleucine 45 µg; L-Arginine hydrochloride 25 µg; L-Phenylalanine 40 µg; L-Valine 50 µg; L-Lysine hydrochloride 50 µg; L-Leucine 40 µg; Monosodium glutamate 50 µg; Thiamine hydrochloride 0.10 mg; Riboflavin 40 µg; Pyridoxine hydrochloride 0.10 mg; Nicotinamide 1.50 mg; D-Panthenol 50 µg, Excipients q.s.1 ml	Bovine Equine	Slow intravenous, intramuscular or subcutaneous routes Calves and Foals: 10 ml/kg b.w./day. Adult Bovines and Equines: 500 - 2000 ml/animal/day, according to weight.	250 ml 500 ml
R-COMPLEX-B Injectable solution	Vitamin B1 hydrochloride 100 mg, Vitamin B2 sodium phosphate 5 mg, Vitamin B6 hydrochloride 50 mg, Vitamin B12 0.10 mg, Nicotinamide 150 mg, D - Panthenol 5 mg, Excipients q.s. 1 ml	Bovine Equine Porcine Ovine Caprine	Intramuscular route Bovine and equine: 1 ml/100 kg b.w./day. Porcine, ovine and caprine: 0.5 ml/50 kg b.w./day. Administer for 2 to 3 days if required.	20 ml 100 ml 250 ml
REGULADOR ORGÁNICO REVEFOS Injectable solution	Vitamin D2 20000 IU; Vitamin E 1 mg; Vitamin B12 50 µg; L-Lysine hydrochloride 10 mg; Arginine aspartate 2 mg; L-Glycine 4.2 mg; DL-Methionine 2.1 mg; Magnesium hypophosphite 2.1 mg; Calcium gluconate 1.5 mg; Sodium chloride 4.5 mg; Potassium iodide 0.15 mg; Calcium hypophosphite 6.25 mg; Excipient q.s. 1 ml	Bovine Equine Ovine Caprine	Intramuscular or subcutaneous route Bovines and adult horses: 10 ml/animal. Calves and foals: 5 ml/animal. Sheep and caprine: 5 ml/animal. Lambs and kids: 2 to 3 ml/animal.	250 ml 500 ml

ANTI-INFLAMMATORY				
FLUNIVEEX Injectable solution	Flunixin (meglumine) 50 mg Excipients q.s. 1 ml	Bovine Equine Porcine	Deep intramuscular or intravenous route Cattle: 2 ml/50 kg b.w. Horses: 2 ml/50 kg b.w. Pigs: 2 ml/50 kg b.w. Administer for no longer than 5 days.	20 ml 50 ml 100 ml
KETOPROFEN Injectable solution	Ketoprofen 100 mg Excipients q.s. 1 ml	Bovine Equine Porcine	Intramuscular or intravenous route Cattle: 0.6 ml/33 kg b.w./day for 1 to 3 days. Horses: 1 ml/45 kg b.w./day for 3 to 5 days. Pigs: 0.6 ml/33 kg b.w./day for 1 to 3 days.	100 ml 250 ml
MELOXICAM 4% Injectable solution	Meloxicam 40 mg Excipients q.s. 1 ml	Cattle Pigs	Horses Intramuscular, SC and IV route. Cattle: 2.5 ml/100 kg b.w. via SC or IV route. Pigs: 2 ml/100 kg b.w. IM. Horses: 3 ml/100 kg b.w. IV.	50 ml 100 ml 250 ml
NOVALGEX Injectable solution	Dipyron 500 mg Excipients q.s. 1 ml	Equine Bovine Ovine Caprine Porcine Canine	Intramuscular or slow intravenous route Adult equines: 20 to 60 ml/animal, depending on weight. Adult cattle: 20 to 40 ml/animal. Foals and calves: 5 to 15 ml/animal. Sheep and goats: 2 to 8 ml/animal. Pigs: 10 to 30 ml/animal. Canines: 1 to 5 ml/animal. Repeat multiple times a day (every 6 to 8 hours).	20 ml 100 ml
RETARDEX Injectable solution	Dexamethasone (21-phosphate sodium) 0.50 mg Dexamethasone (21-isonicotinate) 1 mg Excipient q.s. 1 ml	Bovine Equine Porcine Ovine Caprine	Intramuscular, subcutaneous or intravenous route Bovines and adult horses: 10 ml/animal. Calves, foals, sheep and caprines: 3 to 5 ml/animal Pigs: 2 to 5 ml/animal. Piglets, lambs and kids: 1 to 2 ml/animal.	10 ml 20 ml 50 ml
TOLFEEX Injectable solution	Tolfenamic acid 40 mg Excipients q.s. 1 ml	Bovine Porcine	Intramuscular route Bovines: for inflammation caused by respiratory diseases, use 1 ml/10 kg body weight intramuscularly in the neck muscles. For mastitis, use 1 ml/10 kg body weight intravenously, single dose. Pigs: 1 ml/20 kg body weight in a single intramuscular injection.	20 ml 50 ml 100 ml 250 ml